

- independence

① Prove: If A is independent of B then A is independent of B'.

$$P(A \cap B) = P(A)P(B) \quad P(A) = P(A \cap B) + P(A \cap B')$$

$$= P(A)P(B) + P(A \cap B')$$

$$\Rightarrow P(A)(1 - P(B)) = P(A)P(B') = P(A \cap B')$$

② $E_1, \dots, E_n$ are independent if for every subset $E_{i_1}, \dots, E_{i_r}$ ($r \leq n$)

$$P(E_{i_1} \cap \dots \cap E_{i_r}) = P(E_{i_1}) \dots P(E_{i_r})$$

$$\Rightarrow P(E_1 | f(E_2, \dots, E_n)) = \frac{P(E_1)P(f(E_2, \dots, E_n))}{P(f(E_2, \dots, E_n))} = P(E_1)$$

$f(E_2, \dots, E_n)$ is an event obtained by performing basic operations on sets $E_2, \dots, E_n$.

pairwise independent 两两独立

mutually independent 相互独立

$$\forall i_1, i_2, \dots, i_k \subseteq 1, 2, \dots, n \quad P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = P(A_{i_1})P(A_{i_2}) \dots P(A_{i_k})$$

conditionally independent 条件独立 在已知某个特定信息后, 两事件无关联

$$P(A|C) \cdot P(B|C) = P(A \cap B|C)$$

- the law of total probability

$$P(A) = \sum_{i=1}^n P(A \cap B_i) = \sum_{i=1}^n P(A|B_i)P(B_i)$$

- Bayes' rule

$$P(B_j|A) = \frac{P(A \cap B_j)}{P(A)} = \frac{P(A|B_j) \cdot P(B_j)}{\sum_{i=1}^n P(A|B_i)P(B_i)}$$

[e.g.] C : you have cancer T : you test positive

$$P(T|C) = 0.95 \quad P(T'|C') = 0.95 \quad P(C) = 0.004$$

compute $P(T)$, $P(C|T)$

$$P(T) = P(T|C) \cdot P(C) + P(T|C') \cdot P(C')$$

$$= P(T|C) \cdot P(C) + (1 - P(T'|C')) \cdot (1 - P(C)) = 0.0536$$

$$P(C|T) = \frac{P(C \cap T)}{P(T)} = \frac{P(C) \cdot P(T|C)}{P(T)} = 0.07089$$

• the r -th moment [矩]

1. 矩估计

	样本 (A_r)	总体
① $E(X^r)$ r 阶原点矩	$\frac{1}{n} \sum_{i=1}^n x_i$ $\frac{1}{n} \sum_{i=1}^n x_i^r$	$E(X)$ $E(X^r)$
② $E[(X-b)^r]$ r 阶中心矩	$\frac{1}{n} \sum_{i=1}^n (x_i - b)^r$	$E[(X - E(X))^r]$ ← $r=2 \quad E[(X - E(X))^2] = D[X]$
③ $E[(X)_r] \triangleq E[X(X-1)\cdots(X-r+1)]$ r -th factorial moment		

2. moment generating function [矩母函数]

X be a discrete random variable. if there exists a $h > 0$ s.t.

$$E[e^{tx}] = \sum_{i \in \mathbb{N}} e^{tx_i} f(x_i) \quad \text{exists } \forall t \in (-h, h)$$

$\Rightarrow M(t) = E[e^{tx}]$, $t \in (-h, h)$ is the mgf of X

(连续型: $E[e^{tx}] = \int_{-\infty}^{\infty} f(x) e^{tx} dx$)

$$M'(t) = \frac{dE(e^{tx})}{dt} = E[e^{tx} \cdot x] \quad M'(0) = E(X)$$

$$M^{(r)}(t) = E[X^r e^{tx}] \quad M^{(r)}(0) = E[X^r]$$

$$\text{Var}[X] = M''(0) - [M'(0)]^2$$

e.g.] $X \sim U(a, b)$ (a, b 未知), $X_1, \dots, X_n$ 为来自总体的简单随机样本. 试用矩估计法估计 a, b .

$$\textcircled{1} \bar{X} = EX = \frac{a+b}{2}$$

$$\textcircled{2} \begin{cases} A_2 = \frac{1}{n} \sum_{i=1}^n x_i^2 = E(X^2) = \int_a^b x^2 \frac{1}{b-a} dx = \frac{a^2 + ab + b^2}{3} \\ B_2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 = DX = \frac{(b-a)^2}{12} \quad (\text{better}) \end{cases}$$

$$\Rightarrow \begin{cases} \hat{a} = \bar{x} - \sqrt{3B_2} \\ \hat{b} = \bar{x} + \sqrt{3B_2} \end{cases} \quad \text{* 由于 } B_2 \text{ 可由 } X_1, \dots, X_n \text{ 推出, 故简写为 } B_2$$

3. X is a nonnegative random variable and $c > 0$, then $P(X \geq c) \leq \frac{E[X]}{c}$

Proof:

$$E[X] = \sum_{k \geq 0} k f(k) \geq \sum_{k \geq c} c f(k) \geq c \sum_{k \geq c} f(k) = c P(X \geq c)$$

$$\Rightarrow \frac{E[X]}{c} \geq P(X \geq c)$$

• Distribution

1. Standardized random variable

$$Z = \frac{X - \mu}{\sigma} \quad \begin{cases} E[Z] = \frac{1}{\sigma} [E(X) - \mu] = \frac{1}{\sigma} (\mu - \mu) = 0 \\ \text{Var}(Z) = E[(Z - E(Z))^2] = E[Z^2] = E\left[\frac{1}{\sigma^2} (X - \mu)^2\right] \\ = \frac{1}{\sigma^2} E[(X - \mu)^2] = \frac{1}{\sigma^2} \cdot \sigma^2 = 1 \end{cases}$$

2. Bernoulli Distribution $f(x) = p^x (1-p)^{1-x} \quad x \in \{0, 1\}$

$$E[X] = 1 \cdot p + 0 \cdot (1-p) = p \quad \text{Var}[X] = (1-p)^2 \cdot p + (0-p)^2 (1-p) = p(1-p)$$

$$M(t) = E[e^{tx}] = e^t \cdot p + (1-p) \quad t \in \mathbb{R}$$

3. Binomial Distribution $f(X=k) = \binom{n}{k} p^k (1-p)^{n-k}$

$$E[X] = \sum_{k=0}^n k P(X=k) = np \quad \text{Var}[X] = np(1-p)$$

$$M(t) = \sum_{k=0}^n e^{tk} \cdot \binom{n}{k} p^k (1-p)^{n-k} = [pe^t + (1-p)]^n$$

$$E[X] = M'(0) = np \quad E[X^2] = n(n-1)p^2 + np \quad \text{Var}[X] = np(1-p)$$

4. Hypergeometric Distribution $f(x) = \frac{\binom{N_1}{x} \binom{N_2}{n-x}}{\binom{N}{n}} \quad N = N_1 + N_2$

$$E[X] = n \frac{N_1}{N} \quad \text{Var}[X] = n \frac{N_1}{N} \cdot \frac{N_2}{N} \cdot \frac{N-n}{N-1}$$

when $N \gg n$, $p = \frac{N_1}{N}$, then $E[X] = np$ $\text{Var}[X] = np(1-p)$

5. Geometric Distribution

$$f(x) = (1-p)^{x-1} \cdot p$$

$$E[X] = \sum_{x=1}^{\infty} (1-p)^{x-1} \cdot x \cdot p = \frac{1}{p}$$

$$\text{Var}[X] = \frac{1-p}{p^2}$$

$E[X]$	1	$E(X)+1$
$P(X)$	p	$1-p$

$$M(t) = \frac{pe^t}{1 - (1-p)e^t} \quad (t < -\ln(1-p))$$

$$E[X] = p + (1-p)(E[X]+1) \Rightarrow E[X] = \frac{1}{p}$$

* Negative Binomial Distribution

X : the trial number at which the r -th success is observed.

$r=1$ 几何分布

$$P(X=x) = \binom{x-1}{r-1} \cdot p^{r-1} \cdot (1-p)^{(x-1)-(r-1)} \cdot p = \binom{x-1}{r-1} p^r (1-p)^{x-r} \quad (x \geq r)$$

$$E[X_r] = p \cdot (1 + E[X_{r-1}]) + (1-p) (1 + E[X_r])$$

$$E[X_r] = \frac{1}{p} + E[X_{r-1}] \Rightarrow E[X_r] = \frac{r}{p}$$

$$\text{Var}[X] = \frac{r(1-p)}{p^2}$$

$$M(t) = E[e^{tx}] = \sum_{x=r}^{\infty} e^{tx} \binom{x-1}{r-1} p^r (1-p)^{x-r}$$

$$= p^r e^{tr} \sum_{x=r}^{\infty} \binom{x-1}{r-1} [e^t(1-p)]^{x-r}$$

$$= \frac{(pe^t)^r}{[1 - e^t(1-p)]^r} \quad [\text{if } |e^t(1-p)| < 1, \text{ i.e. } \underline{t < -\ln(1-p)}]$$

$$* (1-w)^{-r} = [1+(-w)]^r = \sum_{k=0}^{\infty} \binom{-r}{k} \cdot (-w)^k$$

$$= \sum_{k=0}^{\infty} (-1)^k \cdot \binom{r+k-1}{k} \cdot (-1)^k \cdot w^k$$

$$= \sum_{k=0}^{\infty} \binom{r+k-1}{k} w^k$$

$$= \sum_{x=r}^{\infty} \binom{x-1}{x-r} w^{x-r} = \sum_{x=r}^{\infty} \binom{x-1}{r-1} w^{x-r}$$

6. Poisson Distribution ① $[0, t_1]$ $[t_1, t_2]$ 事件独立 ② $\lim_{h \rightarrow 0} \frac{P(1 \text{ event in } h)}{h} = \lambda$

$$P(X=x) = e^{-\lambda} \frac{\lambda^x}{x!}$$

③ $\lim_{h \rightarrow 0} \frac{P(\geq 2 \text{ events in } h)}{h} = 0$

$$p = \frac{\lambda}{n} \quad P(X=x) = \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x} \quad n \rightarrow \infty$$

$$\lim_{n \rightarrow \infty} P(X=x) = \frac{n!}{x!(n-x)!} \cdot \left(\frac{\lambda}{n}\right)^x \cdot \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-x}$$

$$\lim_{n \rightarrow \infty} = \frac{n^x}{x!} \cdot \frac{\lambda^x}{n^x} \cdot e^{-\lambda} \cdot 1 = e^{-\lambda} \cdot \frac{\lambda^x}{x!}$$

$$M(t) = \sum_{x=0}^{\infty} e^{tx} e^{-\lambda} \frac{\lambda^x}{x!}$$

$$= e^{-\lambda} \sum_{x=0}^{\infty} \frac{(e^t \lambda)^x}{x!} = e^{-\lambda} \cdot e^{e^t \lambda} = \boxed{e^{\lambda(e^t - 1)}}$$

$$E[X] = M'(0) = \lambda \quad \text{Var}[X] = E[X^2] - (E[X])^2 = \lambda$$

7. Uniform Distribution $E[X] = \frac{1}{2}$, $\text{Var}[X] = \frac{1}{12}$ $M(t) = \begin{cases} \frac{e^t - 1}{t} & t \neq 0 \\ 1 & t = 0 \end{cases}$

X is uniform $[0, 1] \Rightarrow Y = a + (b-a)X$ is uniform $[a, b]$

$$\text{pdf} = \frac{d}{dy} f_Y(y) = \frac{d}{dy} \left(\frac{y-a}{b-a} \right) = \frac{1}{b-a} \quad (a < y < b)$$

$$M_Y(t) = e^{at} M_X((b-a)t)$$

$$E[Y] = a + \frac{1}{2}(b-a) = \frac{a+b}{2}$$

$$\text{Var}[Y] = (b-a)^2 \text{Var}[X] = \frac{(b-a)^2}{12}$$

$$M(t) = \int_a^b e^{ty} \frac{1}{b-a} dy = \frac{1}{b-a} \cdot \frac{1}{t} (e^{tb} - e^{ta}) = \begin{cases} \frac{e^{tb} - e^{ta}}{t(b-a)} & t \neq 0 \\ 1 & t = 0 \end{cases}$$

8. 100th Percentile

$$p = \int_{-\infty}^{\pi_p} f(x) dx = F(\pi_p)$$

[e.g.] $Y \sim U(a, b)$

$$p = \int_{-\infty}^{\pi_p} f(x) dx = \int_{-\infty}^{\pi_p} \frac{1}{b-a} dx = \frac{\pi_p - a}{b-a} \Rightarrow \pi_p = p(b-a) + a$$

9. Exponential Distribution

指数分布 事件的时间间隔概率 [何时]

$$f(x) = \frac{1}{\theta} e^{-\frac{x}{\theta}} \quad (0 \leq x < \infty, \theta > 0)$$

伯松分布 某段时间内, 事件具体发生的概率 [次数]

$$\text{CDF: } F(x) = \begin{cases} 0 & -\infty < x < 0 \\ 1 - e^{-\frac{x}{\theta}} & 0 \leq x < \infty \end{cases}$$

$$P(X > k) = \int_k^{\infty} \frac{1}{\theta} e^{-\frac{x}{\theta}} dx = e^{-\frac{k}{\theta}}$$

$$E[X] = \theta, \text{Var}[X] = \theta^2$$

$$M(t) = \int_0^{\infty} e^{tx} \frac{1}{\theta} e^{-\frac{x}{\theta}} dx = \boxed{\frac{1}{1-t\theta}} \quad (t < \frac{1}{\theta})$$

$$P(X > t+s | X > s) = \frac{P(X > t+s)}{P(X > s)} = e^{-\frac{t}{\theta}} = P(X > t) \quad \text{无记忆性}$$

$$[\text{几何分布 } P(X > i+j | X > i) = \frac{(1-p)^{i+j}}{(1-p)^i} = (1-p)^j = P(X > j)]$$

$$\underline{P(X > k) = (1-p)^k}$$

[e.g.] Customer arrive in a certain shop according to a poisson process at a mean rate of 20 per hour. P (the shopkeeper will have to wait more than 5 minutes for the arrival of the first customer?)

$$\lambda = \frac{1}{3} \text{ person per minute 发生率 } \theta = \frac{1}{\lambda} = 3$$

$$\Rightarrow f(x) = \frac{1}{3} e^{-\frac{x}{3}} (x \geq 0) \quad F(x) = 1 - e^{-\frac{x}{3}} \quad \Rightarrow P(X > 5) = 1 - F(5) \approx 0.189$$

10. Erlang Distribution [第 α 次发生所需时间]

$$\text{CDF} \quad F_W(t) = 1 - P(W > t) = 1 - \sum_{k=0}^{\alpha-1} e^{-\lambda t} \frac{\lambda^k t^k}{k!}$$

$\underbrace{\quad}_{P(Y < \alpha)}$

$$\text{PDF} \quad f_W(t) = f'_W(t) = \frac{\lambda^\alpha}{(\alpha-1)!} t^{\alpha-1} e^{-\lambda t} \quad (t \geq 0)$$

$$\text{平均时间间隔} = \frac{1}{\lambda} \downarrow = \frac{1}{\theta} \quad = \frac{1}{(\alpha-1)! \theta^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta}} \quad (0 \leq x < \infty) \quad \theta = \frac{1}{\lambda}$$

$$E[W] = \alpha \theta \quad \text{Var}[W] = \alpha \theta^2$$

11. Gamma Distribution $\alpha \in \mathbb{Z} \Rightarrow \alpha \in \mathbb{R}$

$$f(x) = \frac{1}{\Gamma(\alpha) \theta^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta}}$$

$$\frac{1}{\Gamma(\alpha)} \int_0^\infty \frac{1}{\theta^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta}} dx = 1 \Rightarrow \Gamma(\alpha) = \int_0^\infty \frac{1}{\theta^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta}} dx$$

$$= \int_0^\infty y^{\alpha-1} e^{-y} dy$$

$$\textcircled{1} \Gamma(\alpha) = (\alpha-1) \Gamma(\alpha-1)$$

$$\Gamma(\alpha) = [-y^{\alpha-1} e^{-y}]_0^\infty + \int_0^\infty (\alpha-1) y^{\alpha-2} e^{-y} dy = (\alpha-1) \int_0^\infty y^{\alpha-2} e^{-y} dy = (\alpha-1) \Gamma(\alpha-1)$$

$$\textcircled{2} \alpha = n. \Gamma(\alpha) = (n-1)!$$

$$\Gamma(n) = (n-1) \Gamma(n-1) \dots = (n-1)(n-2) \dots 1 \cdot \Gamma(1) = (n-1)!$$

$$\Gamma(1) = \int_0^\infty e^{-y} dy = -e^{-y} \Big|_0^\infty = 1$$

$$M(t) = E[e^{tx}] = \int_0^{\infty} e^{tx} \cdot \frac{1}{\Gamma(\alpha)\theta^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta}} dx = \frac{1}{\Gamma(\alpha)\theta^\alpha} \int_0^{\infty} x^{\alpha-1} \exp[-(\frac{1}{\theta}-t)x] dx$$

$$\text{let } \beta = \frac{1}{\theta} - t, \quad M(t) = \frac{1}{\Gamma(\alpha)\theta^\alpha} \int_0^{\infty} x^{\alpha-1} e^{-\beta x} dx \quad (t < \frac{1}{\theta})$$

$$\Gamma(\alpha) = \int_0^{\infty} y^{\alpha-1} e^{-y} dy, \quad M(t) = \frac{1}{\Gamma(\alpha)\theta^\alpha} \cdot \int_0^{\infty} (\frac{y}{\beta})^{\alpha-1} e^{-y} (\frac{dy}{\beta}) = \frac{1}{\beta^\alpha} \cdot \frac{1}{\Gamma(\alpha)\theta^\alpha} \cdot \Gamma(\alpha)$$

let $y = \beta x$ ($x, y \in (0, \infty)$)

$$= \frac{1}{(\theta\beta)^\alpha} = \frac{1}{(1-\theta t)^\alpha}$$

$$E[X] = \alpha\theta, \quad \text{Var}[X] = \alpha(\alpha+1)\theta^2 - [\alpha\theta]^2 = \alpha\theta^2 \quad E[Y^k] = \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)} \theta^k = \frac{(\alpha+k-1)!}{(\alpha-1)!} \theta^k$$

12. Chi-square Distribution 卡方分布 $X \sim \chi^2(r)$

$$f(x) = \frac{1}{\Gamma(\frac{r}{2})2^{\frac{r}{2}}} x^{\frac{r}{2}-1} e^{-\frac{x}{2}} \quad 0 < x < \infty \quad [\text{在Gamma分布中取 } \alpha = \frac{r}{2}, \theta = 2]$$

$$M(t) = (1-2t)^{-\frac{r}{2}} \quad \mu = \alpha\theta = r \quad \sigma^2 = \alpha\theta^2 = 2r$$

$$X_1 \sim X_n \text{ 相互独立, } X_i \sim N(0,1) \quad X = \sum_{i=1}^n X_i^2 \sim \chi^2(n) \quad n: \text{degrees of freedom}$$

$$E[X^2] = n, \quad D(X^2) = 2n$$

13. Normal Distribution $X \sim N(\mu, \sigma^2)$

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) \quad x \in \mathbb{R}$$

$$M(t) = \int_{-\infty}^{\infty} \frac{e^{tx}}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) dx$$

$$= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{1}{2\sigma^2}(x^2 - 2(\mu + \sigma^2 t)x + \mu^2)\right] dx$$

$$= \exp\left(\frac{2\mu\sigma^2 t + \sigma^4 t^2}{2\sigma^2}\right) \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{[x - (\mu + \sigma^2 t)]^2}{2\sigma^2}\right] dx$$

$X \sim N(\mu + \sigma^2 t, \sigma^2)$

$$= \exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right)$$

$$E[X] = M'(0) = \mu, \quad \text{Var}[X] = (\mu^2 + \sigma^2) - \mu^2 = \sigma^2$$

* Standard normal distribution $X \sim N(0,1)$

$$Z_\alpha \Rightarrow P(Z \geq Z_\alpha) = \alpha \quad Z_\alpha = \Phi^{-1}(1-\alpha) \quad 100(1-\alpha)\text{-th percentile}$$

$X \sim N(u, \sigma^2)$. then $Z = \frac{x-u}{\sigma}$ is $N(0,1)$

Proof:

$$P(Z \leq z) = P\left(\frac{x-u}{\sigma} \leq z\right) = P(x \leq z\sigma + u) = \int_{-\infty}^{z\sigma+u} \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{(x-u)^2}{2\sigma^2}\right] dx$$

$$\text{let } w = \frac{x-u}{\sigma}, \text{ then } P(Z \leq z) = \int_{-\infty}^z \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{w^2}{2}} \cdot \sigma dw \\ = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} e^{-\frac{w^2}{2}} dw \Rightarrow u' = 0, \sigma^2 = 1$$

$$* P(a \leq X \leq b) = P\left(\frac{a-u}{\sigma} \leq \frac{x-u}{\sigma} \leq \frac{b-u}{\sigma}\right) = \Phi\left(\frac{b-u}{\sigma}\right) - \Phi\left(\frac{a-u}{\sigma}\right)$$

* Normal and χ^2 Distribution

$$X \sim N(u, \sigma^2) \quad V = \frac{(X-u)^2}{\sigma^2} = Z^2 \text{ is } \chi^2(1) \quad V \sim \chi^2(1)$$

Proof:

$$G(v) : \text{the cdf} \quad G(v) = P(Z^2 \leq v) = \int_{-\sqrt{v}}^{\sqrt{v}} \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz = \int_0^v \frac{1}{\sqrt{2\pi y}} e^{-\frac{y}{2}} dy \\ g(v) = G'(v) = \frac{1}{\sqrt{2\pi v}} e^{-\frac{v}{2}} \quad (z = \sqrt{y})$$

$$= \frac{1}{\sqrt{\pi} \cdot \sqrt{2}} v^{\frac{1}{2}-1} \cdot e^{-\frac{v}{2}}$$

$$X \sim \chi^2(1) : f(x) = \frac{1}{\Gamma(\frac{1}{2})\sqrt{2}} x^{\frac{1}{2}-1} e^{-\frac{x}{2}}$$

$$\int_0^\infty g(v) dv = 1 \Rightarrow \frac{1}{\sqrt{2\pi}} \int_0^\infty (2x)^{\frac{1}{2}-1} e^{-x} \cdot 2 dx \quad \int_0^\infty x^{\alpha-1} e^{-x} dx \\ = \frac{1}{\sqrt{\pi}} \int_0^\infty x^{\frac{1}{2}-1} \cdot e^{-x} dx = \frac{1}{\sqrt{\pi}} \Gamma(\alpha) = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1}{2}\right) = 1$$

$$\Rightarrow \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

$$\Rightarrow \text{cdf } g(v) \quad \underline{\chi^2(r)} \quad r=1 \text{ 的情况}$$

• Bivariate Distribution of Discrete Type [二维]

1. Marginal Probability mass function 边际概率质量函数

$$f_X(x) = \sum_y f(x, y) = P(X=x)$$

independent $f(x, y) = f_X(x) \cdot f_Y(y)$ ★点、和集合均可证明独立

① x, Y independent $\Rightarrow P(X \in A, Y \in B) = P(X \in A) P(Y \in B)$

$$P(X \in A, Y \in B) = \sum_{y \in B} P(X \in A, Y=y)$$

$$= \sum_{y \in B} \sum_{x \in A} P(X=x, Y=y) = \sum_{y \in B} \sum_{x \in A} P(X=x) \cdot P(Y=y)$$

$$= \sum_{y \in B} P(X \in A) P(Y=y) = P(X \in A) P(Y \in B)$$

② 集合独立 $\Rightarrow$ 点独立

取 $A = \{x\}$, $B = \{y\}$ 利用单点集即可证明

2. Joint Cumulative distribution function

$$F(x, y) = P(X \leq x, Y \leq y) \quad [X, Y \text{ independent} \Leftrightarrow F(x, y) = F_X(x) \cdot F_Y(y)]$$

Lemma: $F(x, y) = F_X(x) \cdot F_Y(y) \Rightarrow \{X=x_i\}, \{Y \leq y\}$ independent $\forall x_i \in S_X, y$

$$P(X=x_i, Y \leq y) = P(X \leq x_i, Y \leq y) - P(X \leq x_{i-1}, Y \leq y) \quad (\text{假设 } x, y \text{ 已升序排列})$$

$$= [P(X \leq x_i) - P(X \leq x_{i-1})] P(Y \leq y) = P(X=x_i) P(Y \leq y) \quad (i=1, x_0=x_{i-1})$$

① $F(x, y) = F_X(x) F_Y(y) \Rightarrow X, Y$ independent

$$P(X=x_i, Y=y_j) = P(X=x_i, Y \leq y_j) - P(X=x_i, Y \leq y_{j-1}) = P(X=x_i) P(Y=y_j)$$

$$\text{i.e. } f(x_i, y_j) = f_X(x_i) f_Y(y_j) \quad (j=1, y_0=y_{j-1})$$

$$F(x, y) = \int_{s=-\infty}^x \int_{t=-\infty}^y f(s, t) dt ds = \int_{s=-\infty}^x \int_{t=-\infty}^y f_X(s) \cdot f_Y(t) dt ds$$

$$= \int_{s=-\infty}^x f_X(s) \int_{t=-\infty}^y f_Y(t) dt ds = F_X(x) F_Y(y)$$

② X, Y independent $\Rightarrow F(x, y) = F_X(x) \cdot F_Y(y)$

$$F(x, y) = P(X \leq x, Y \leq y) = P(X \leq x) P(Y \leq y) = F_X(x) \cdot F_Y(y)$$

$$3. E[X] = \sum_{x \in S_X} x f_X(x) = \sum_{(x,y) \in S} x f(x,y)$$

$$E[u(x_1, x_2)] = \sum_{(x_1, y_1) \in S} \sum_{(x_2, y_2) \in S} u(x_1, x_2) \cdot f(x_1, x_2)$$

4. Covariance

$$\text{COV}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)] = \sigma_{XY}$$

$$= E[XY - \mu_X Y - \mu_Y X + \mu_X \mu_Y]$$

$$= E[XY] - E[X] E[Y]$$

$$\cdot \text{COV}(X, Y) \begin{cases} > 0 & \text{positively correlated} \\ = 0 & \text{uncorrelated} \\ < 0 & \text{negatively correlated} \end{cases}$$

$$\cdot X, Y \text{ independent} \Rightarrow \text{COV}(X, Y) = 0 \quad (\text{反之不一定成立})$$

$$E[XY] = \sum_{(x,y) \in S} xy f(x,y) = \sum_{(x,y) \in S} xy f_X(x) f_Y(y)$$

$$= \sum_{x \in S_X} x f_X(x) \cdot \sum_{y \in S_Y} y f_Y(y) = E[X] E[Y]$$

$$\cdot \text{COV}(X, X) = \text{Var}[X]$$

5. The Correlation Coefficient

$$\rho = \frac{\text{COV}(X, Y)}{\sigma_X \sigma_Y} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$$

$$\textcircled{1} \rho \in [-1, 1] \Leftrightarrow \text{COV}(X, Y)^2 \leq (\sigma_X \sigma_Y)^2$$

$$\text{let } V = X - \mu_X, W = Y - \mu_Y$$

$$E[(V + tW)^2] = E[V^2] + 2tE[VW] + t^2 E[W^2]$$

$$= \sigma_X^2 + 2t \operatorname{Cov}(X, Y) + t^2 \sigma_Y^2 \geq 0 \quad (1)$$

$\forall t \in \mathbb{R}$, (1) is true

$$\Rightarrow t = -\frac{\operatorname{Cov}(X, Y)}{\sigma_Y^2} \quad f(t) = \frac{\operatorname{Cov}^2(X, Y)}{\sigma_Y^2} - \frac{2\operatorname{Cov}^2(X, Y)}{\sigma_Y^2} + \sigma_X^2 \geq 0$$

$$\Rightarrow \operatorname{Cov}(X, Y)^2 \leq (\sigma_X \sigma_Y)^2$$

$$\bullet \rho = \pm 1 \quad Y - E[Y] = K(X - E[X]) \quad \exists K \in \mathbb{R}$$

$$\rho = \pm 1, \operatorname{Cov}(X, Y) = \pm \sigma_X \sigma_Y \quad \text{find } K \Rightarrow \text{find } t \text{ s.t. } E[(V + tW)] = 0$$

$$\text{since } E[(V + tW)^2] \geq 0, \text{ check } f(t)_{\min} = f\left(-\frac{\operatorname{Cov}(X, Y)}{\sigma_Y^2}\right) = 0$$

$$\Rightarrow V + \left[-\frac{\operatorname{Cov}(X, Y)}{\sigma_Y^2}\right]W = 0 \Rightarrow K = \frac{\operatorname{Cov}(X, Y)}{\sigma_Y^2}$$

6. Conditional Probability

$$g(x|y) = P(X=x | Y=y) = \frac{P(A \cap B)}{P(B)} \quad \leftarrow A = \{X=x\}, B = \{Y=y\}$$

$$= \frac{f(x, y)}{f_Y(y)}$$

$$\bullet \text{ if } f(x, y) = f_X(x) f_Y(y) \Rightarrow g(x|y) = f_X(x)$$

$$\sum_{y \in S_Y} h(y|x) = \sum_{y \in S_Y} \frac{f(x, y)}{f_X(x)} = \frac{\sum_{y \in S_Y} f(x, y)}{f_X(x)} = \frac{f_X(x)}{f_X(x)} = 1$$

$$\bullet E[u(x) | Y=y] = \sum_{x \in S_X} u(x) g(x|y)$$

$$= \sum_{x \in S_X} x \cdot g(x|y)$$

$$\bullet \operatorname{Var}[X | Y=y] = \sum_{x \in S_X} (x - E[X | Y=y])^2 g(x|y)$$

[e.g.] X be the number of success in n Bernoulli trials.

Y be the number of success in the first $m < n$ trials.

Z be the number of success in the last $n-m$ trials.

$[Z \text{ is independent of } Y]$

$$\textcircled{1} g(x|y) = \frac{f(x,y)}{f_Y(y)} = \frac{f(x-y,y)}{f_Y(y)} \leftarrow x = y+z$$

$$\begin{matrix} z \in [0, n-m] \\ \downarrow \\ n-m+y \end{matrix} = \frac{f_Y(y) f_Z(x-y)}{f_Y(y)} = f_Z(x-y)$$

$$\textcircled{2} E[X|Y=y] = \sum_{x=y} x f_Z(x-y) \quad Z \sim B(n-m, p)$$

$$= \sum_{z=0}^{n-m} (y+z) f_Z(z)$$

$$= y \sum_{z=0}^{n-m} f_Z(z) + \sum_{z=0}^{n-m} z f_Z(z)$$

$$= y \cdot 1 + E[Z] = y + (n-m)p$$

$$\textcircled{3} \text{Var}[X|Y=y] = \sum_{x=y}^{n-m+y} [x - y - (n-m)p]^2 f_Z(x-y)$$

$$= \sum_{z=0}^{n-m} [z - (n-m)p]^2 f_Z(z) = \text{Var}[Z]$$

$$= (n-m)p(1-p)$$

7. Joint Probability Density Function

$$P((X,Y) \in A) = \iint_A f(x,y) dx dy \quad A \subset \mathbb{R}^2$$

$$E[u(X,Y)] = \iint_{(x,y) \in \mathbb{S}} u(x,y) f(x,y) dx dy$$

$$\text{Var}[X] = E[(X - E[X])^2]$$

$$\text{Var}[Y] = E[(Y - E[Y])^2]$$

$$\text{Cov}(X,Y) = E[XY] - E[X]E[Y]$$

$$E[XY] = \iint_{(x,y) \in \mathbb{S}} xy f(x,y) dx dy, \quad \rho(X,Y) = \frac{\text{Cov}(X,Y)}{\sqrt{\text{Var}[X]\text{Var}[Y]}}$$

[e.g.] $f(x,y) = \frac{1}{\theta^2} \exp(-\frac{y}{\theta}) \quad 0 < x < y < \infty$

① $E[X], \text{Var}[X], E[X|Y=y], \text{Var}[X|Y=y]$

② $E[Y], \text{Var}[Y], E[Y|X=x], \text{Var}[Y|X=x]$

$$f_X(x) = \int_{y=x}^{\infty} f(x,y) dy = \theta e^{-\frac{x}{\theta}} \quad X \sim \text{Exp}(\theta) \Rightarrow E[X] = \theta, \text{Var}[X] = \theta^2$$

$$f_Y(y) = \int_{x=0}^y f(x,y) dx = \frac{y}{\theta^2} e^{-\frac{y}{\theta}} \quad Y \sim \text{lang}(\alpha=2) \Rightarrow E[Y] = 2\theta, \text{Var}[Y] = 2\theta^2$$

$$f_X|Y(X|Y) = \frac{f(X,Y)}{f_Y(Y)} = \frac{1}{Y} \quad X|Y=y \sim (0, Y) \rightarrow E[X|Y=y] = \frac{Y}{2} \quad \text{Var}[X|Y=y] = \frac{Y^2}{12}$$

$$f_Y|X(Y|X) = \frac{1}{\theta} e^{-\frac{Y-X}{\theta}} \quad E[Y|X=x] = E[X+Z] = \theta + x$$

$$Y = X + Z \quad \text{Var}[Y|X=x] = \text{Var}[X+Z] = \text{Var}[Z] = \theta^2$$

constant

8. tower Property $E[E[X|Y]] = E[X]$

proof: $E[E[X|Y]] = \int_{S_Y} E[X|Y=y] f_Y(y) dy = \int_{S_Y} \int_R x g(x|y) f_Y(y) dx dy$

$$= \int_R x \int_{S_Y} f(x,y) dy dx$$

$$= \int_R x f_X(x) dx = E[X]$$

$$\text{Var}[Y] = E[\text{Var}[Y|X]] + \text{Var}[E[Y|X]]$$

$$E[Y^2] = E[E[Y^2|X]] = E[\text{Var}[Y|X] + (E[Y|X])^2] = E[\text{Var}[Y|X]] + E[E[Y|X]^2]$$

$$\text{Var}[Y] = E[Y^2] - (E[Y])^2 = E[\text{Var}[Y|X]] + E[E[Y|X]^2] - (E[Y])^2$$

$$= E[\text{Var}[Y|X]] + E[E[Y|X]^2] - (E[E[Y|X]])^2$$

$$= E[\text{Var}[Y|X]] + \text{Var}[E[Y|X]]$$

9. Bivariate Normal Distribution

$$f(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left[-\frac{q(x,y)}{2}\right], \quad q(x,y) = \frac{1}{1-\rho^2} \left[\left(\frac{x-\mu_X}{\sigma_X}\right)^2 - 2\rho \left(\frac{x-\mu_X}{\sigma_X}\right) \left(\frac{y-\mu_Y}{\sigma_Y}\right) + \left(\frac{y-\mu_Y}{\sigma_Y}\right)^2 \right]$$

$\sigma_X > 0, \sigma_Y > 0, |\rho| < 1$

* 边缘分布均为正态, $X \sim N(\mu_X, \sigma_X^2), Y \sim N(\mu_Y, \sigma_Y^2)$

$$E[X|Y=y] = \mu_X + \frac{\sigma_X}{\sigma_Y} \rho (y - \mu_Y) \quad \text{Var}[X|Y=y] = (1-\rho^2) \sigma_X^2$$

$$E[Y|X=x] = \mu_Y + \frac{\sigma_Y}{\sigma_X} \rho (x - \mu_X)$$

$$\star E[XY] = E[E[XY|X]]$$

$$= E[X \cdot E[Y|X]]$$

$\star$ independence $\Leftrightarrow$ uncorrelation

proof:

$$\textcircled{1} f_X(x) = \int_{-\infty}^{\infty} f(x,y) dy = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(-\frac{u^2}{2}\right) \int_{-\infty}^{\infty} \exp\left[-\frac{(v-\rho u)^2}{2(1-\rho^2)}\right] dv$$

$u = \frac{x-\mu_X}{\sigma_X}, v = \frac{y-\mu_Y}{\sigma_Y} \Rightarrow q(x,y) = \frac{1}{1-\rho^2} (u^2 - 2\rho uv + v^2)$

$$= \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(-\frac{u^2}{2}\right) \int_{-\infty}^{\infty} \exp\left[-\frac{t^2}{2(1-\rho^2)}\right] dt \quad (t = v - \rho u) = \frac{1}{\sqrt{2\pi}\sigma_X} \exp\left[-\frac{(x-\mu_X)^2}{2\sigma_X^2}\right]$$

$$\textcircled{2} g(x|y) = \frac{f(x,y)}{f_Y(y)} = \frac{\frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left[-\frac{q(x,y)}{2}\right]}{\frac{1}{\sqrt{2\pi}\sigma_Y} \exp\left[-\frac{(y-\mu_Y)^2}{2\sigma_Y^2}\right]} = \frac{1}{\sqrt{2\pi}\sigma_X\sqrt{1-\rho^2}} \exp\left[-\frac{(x-B)^2}{2(1-\rho^2)\sigma_X^2}\right]$$

$B = \mu_X + \rho \frac{\sigma_X}{\sigma_Y} (y - \mu_Y)$

$$\Rightarrow E[X|Y=y] = \mu_X + \frac{\sigma_X}{\sigma_Y} \rho (y - \mu_Y) \quad \text{Var}[X|Y=y] = (1-\rho^2) \sigma_X^2$$

$$\textcircled{3} 1^\circ \text{ independent} \Rightarrow \rho = \frac{\text{COV}(X,Y)}{\sigma_X\sigma_Y} = 0$$

$$2^\circ \rho = 0 \quad f(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y} \exp\left[-\frac{1}{2} \left(\frac{(x-\mu_X)^2}{\sigma_X^2} + \frac{(y-\mu_Y)^2}{\sigma_Y^2} \right)\right]$$

$$= \frac{1}{\sqrt{2\pi}\sigma_X} \exp\left[-\frac{(x-\mu_X)^2}{2\sigma_X^2}\right] \cdot \left[\frac{1}{\sqrt{2\pi}\sigma_Y} \exp\left[-\frac{(y-\mu_Y)^2}{2\sigma_Y^2}\right] \right] = f_X(x) \cdot f_Y(y)$$

10. function of one random variable

[one-to-one mapping]

$$Y = u(X) \Leftrightarrow X = v(Y)$$

<1> discrete case

$$g(y) = P(Y=y) = P(u(X)=y) = P(X=v(y))$$

$$P(X=x) = f(x) \Rightarrow g(y) = f(v(y))$$

<2> continuous case $\rightarrow$ 反函数唯一

[monotonic function] $g(y) = f(v(y)) \left| \frac{dv(y)}{dy} \right|$

① increasing

$$G(y) = P(Y \leq y) = P(u(x) \leq y) = P(x \leq v(y)) = \int_{c_1}^{v(y)} f(x) dx$$

$$g(y) = G'(y) = f(v(y)) v'(y) = f(v(y)) \left| \frac{dv(y)}{dy} \right|$$

② decrease $G(y) = P(x \geq v(y)) = 1 - \int_{c_1}^{v(y)} f(x) dx$

$$g(y) = -f(v(y)) \frac{dv(y)}{dy} = f(v(y)) \left| \frac{dv(y)}{dy} \right|$$

• random number generator

$f(x)$ strictly increasing

<1> $Y \sim U(0,1)$ $a < x < b$ ($f(a)=0, f(b)=1$) $\Rightarrow X = f^{-1}(Y)$ is a

continuous-type random variable with cdf $f(x)$

proof: $P(X \leq x) = P(f^{-1}(Y) \leq x)$ $\downarrow$ 单调递增

$$= P(f(f^{-1}(Y)) \leq f(x)) = P(Y \leq f(x))$$

$$Y \sim U(0,1) \Rightarrow P(X \leq x) = P(Y \leq f(x)) = f(x)$$

<2> cdf $f(x)$ $a < x < b$ strictly increasing

$$Y = f(X) \sim U(0,1)$$

proof: $f(a)=0, f(b)=1 \Rightarrow P(Y \leq y) = P(f(x) \leq y) \quad y \in (0,1)$

$$= P(X \leq f^{-1}(y))$$

$$P(X \leq x) = f(x) \Rightarrow P(X \leq f^{-1}(y)) = f(f^{-1}(y)) = y \sim U(0,1)$$

<3> X random variable $\Rightarrow$ find $Y = v(x)$

[e.g.] $f(x) = \frac{1}{\pi(1+x^2)}$, $x \in (-\infty, \infty)$, $Y = X^2$

find the pdf of Y

$$G(y) = P(Y \leq y) = P(X^2 \leq y) = \int_{-\sqrt{y}}^{\sqrt{y}} f(x) dx = 2 \int_0^{\sqrt{y}} \frac{1}{\pi(1+x^2)} dx$$

$$g(y) = G'(y) = \frac{1}{\pi(1+y)\sqrt{y}}$$

• Several random variables

1. n independent random variables [mutually independent]

$$f(x_1, \dots, x_n) = f_1(x_1) \cdots f_n(x_n)$$

2. Independently and identically distributed random variables

$$g(x_1, \dots, x_n) = f(x_1) \cdots f(x_n) \quad [\text{直接分解成 } n \text{ 个相同的分布}]$$

★ $X_1, \dots, X_n$ independent random variables

$$E[U_1(X_1) \cdots U_n(X_n)] = \sum_{x_1 \in \mathcal{S}_{X_1}} U_1(x_1) f_1(x_1) \cdots \sum_{x_n \in \mathcal{S}_{X_n}} U_n(x_n) f_n(x_n) \\ = E[U_1(X_1)] \cdots E[U_n(X_n)]$$

$$Y = \sum_{i=1}^n a_i X_i$$

$$E[Y] = \sum_{i=1}^n a_i E[X_i] = \sum_{i=1}^n a_i \mu_i$$

$$\text{Var}[Y] = E\left[\left(\sum_{i=1}^n a_i X_i - \sum_{i=1}^n a_i \mu_i\right)^2\right] = E\left[\left(\sum_{i=1}^n a_i (X_i - \mu_i)\right)^2\right] = \sum_{i=1}^n \sum_{j=1}^n a_i a_j E[(X_i - \mu_i)(X_j - \mu_j)] \\ \text{when } i \neq j, \text{ 由独立性和 } E[(X_i - \mu_i)(X_j - \mu_j)] = E[X_i - \mu_i] \cdot E[X_j - \mu_j] = 0 \rightarrow \sum_{i=1}^n a_i^2 \sigma_i^2 \quad (i=j, E[(X_i - \mu_i)^2] = \sigma_i^2)$$

4. Moment Generating function Technique

" $X_1, \dots, X_n$ are independent random variables, $M_{X_i}(t) \quad t \in [-h_i, h_i]$, $Y = \sum_{i=1}^n a_i X_i$ "

$$M_Y(t) = E[e^{tY}] = E[e^{t(a_1 X_1 + \dots + a_n X_n)}] = E[e^{a_1 t X_1}] \cdots E[e^{a_n t X_n}] \quad (\text{by independence}) \\ = \prod_{i=1}^n M_{X_i}(a_i t)$$

$$\Rightarrow M_{X_i}(t) = M(t) \quad \textcircled{1} \quad M_Y(t) = [M(t)]^n \quad (a=1)$$

$$\textcircled{2} \quad \bar{X} = \sum_{i=1}^n \left(\frac{1}{n}\right) X_i \quad M_{\bar{X}}(t) = \left[M\left(\frac{t}{n}\right)\right]^n \quad (a=\frac{1}{n})$$

(2) $X_1, \dots, X_n$ be independent chi-square random variables with $r_1, \dots, r_n$ degrees of freedom $Y = X_1 + X_2 + \dots + X_n \sim \chi^2(r_1 + \dots + r_n)$

$$X_i \sim \chi^2(r_i) \quad M_{X_i}(t) = \frac{1}{(1-2t)^{\frac{r_i}{2}}} \quad t < \frac{1}{2} \quad M_Y(t) = \frac{1}{(1-2t)^{\frac{r_1+r_2+\dots+r_n}{2}}} \quad (t < \frac{1}{2}) \quad \text{i.e. } \sim \chi^2(r_1 + \dots + r_n)$$

$$\Rightarrow X_i \sim N(0, 1) \quad X_i^2 \sim \chi^2(1) \quad Y = \sum X_i^2 \sim \chi^2(n)$$

$$X_i \sim N(\mu_i, \sigma_i^2) \Rightarrow Y = \sum \frac{(X_i - \mu_i)^2}{\sigma_i^2} \sim \chi^2(n)$$

5.

(1) $X_i \sim N(\mu_i, \sigma_i^2)$ mutually independent $Y = \sum_{i=1}^n c_i X_i \sim N\left(\sum_{i=1}^n c_i \mu_i, \sum_{i=1}^n c_i^2 \sigma_i^2\right)$

$$M_Y(t) = \prod_{i=1}^n M_{X_i}(c_i t) = \prod_{i=1}^n \exp\left(\mu_i c_i t + \frac{\sigma_i^2 c_i^2 t^2}{2}\right) = \exp\left[\left(\sum_{i=1}^n c_i \mu_i\right)t + \left(\sum_{i=1}^n c_i^2 \sigma_i^2\right)\frac{t^2}{2}\right]$$

$$* X_i \sim N(\mu, \sigma^2) \Rightarrow \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N\left(\mu, \frac{\sigma^2}{n}\right) \quad [c_i = \frac{1}{n}]$$

$$(2) X_i \sim N(\mu, \sigma^2), \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \quad (2) \frac{(n-1)S^2}{\sigma^2} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2} \sim \chi^2(n-1)$$

无偏方差 Δ

$$\text{Proof (2): } W = \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 = \sum_{i=1}^n \left[\frac{(X_i - \bar{X}) + (\bar{X} - \mu)}{\sigma} \right]^2 = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 + \frac{n(\bar{X} - \mu)^2}{\sigma^2} + 2 \sum_{i=1}^n \frac{(\bar{X} - \mu)(X_i - \bar{X})}{\sigma^2}$$

$$= \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 + \frac{n(\bar{X} - \mu)^2}{\sigma^2} + \frac{2(\bar{X} - \mu)}{\sigma^2} \sum_{i=1}^n (X_i - \bar{X}) = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 + \frac{n(\bar{X} - \mu)^2}{\sigma^2} \sim \chi^2(n)$$

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) \Rightarrow \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \sim N(0, 1) \Rightarrow \frac{n(\bar{X} - \mu)^2}{\sigma^2} \sim \chi^2(1)$$

$$M(t) = \frac{1}{(1-2t)^{\frac{n}{2}}} = M_1(t) \cdot \frac{1}{(1-2t)^{\frac{1}{2}}} \Rightarrow M_1(t) = \frac{1}{(1-2t)^{\frac{n-1}{2}}}, \text{ i.e. } \chi^2(n-1)$$

* $\bar{X}$ 代替 μ $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \Rightarrow \sum_{i=1}^n (X_i - \bar{X}) = 0$ 由 $n-1$ 个离差 $\rightarrow n$ 个约束自动满足

6. Student's t Distribution (学生氏分布) 相较于 N , T 分布容错率更高 $\uparrow$ 更接近正态分布

$$T = \frac{Z}{\sqrt{\frac{U}{r}}}, \quad Z \sim N(0, 1), U \sim \chi^2(r), Z, U \text{ is independent}$$

$$f(t) = \frac{\Gamma(\frac{r+1}{2})}{\sqrt{\pi r} \Gamma(\frac{r}{2})} \frac{1}{(1 + \frac{t^2}{r})^{\frac{r+1}{2}}}, \quad -\infty < t < \infty$$

$$\text{proof: } g(z, u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} \frac{1}{\Gamma(\frac{r}{2}) 2^{\frac{r}{2}}} u^{\frac{r}{2}-1} e^{-\frac{u}{2}}, \quad z \in \mathbb{R}, u \in [0, \infty)$$

$$f(t) = P(T \leq t) = P(Z \leq \sqrt{\frac{U}{r}} t) \quad f(t) = F'(t)$$

$$\uparrow Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \sim N(0, 1) \quad U = \frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$$

$$T = \frac{Z}{\sqrt{\frac{U}{n-1}}} = \frac{\bar{X} - \mu}{\frac{s}{\sqrt{n}}} \sim t(n-1) \quad \leftarrow \begin{array}{l} \text{总体标准差} \\ \downarrow \\ \text{未知 } \sigma, \text{ 用 } S \text{ 估计} \end{array}$$

[$n \geq 30$, by LCT $\rightarrow$ normal distribution]

7. Central Limit Theorem

① $\lim_{n \rightarrow \infty} f_n(z) = f(z)$ (cont.) f 是 cdf, 则有 $Z_n \xrightarrow{d} z$ 依分布收敛

② CLT: $X_1, \dots, X_n \sim N(\mu, \sigma^2), \sigma^2 > 0$

$$\frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \xrightarrow{d} N(0, 1) \Rightarrow \begin{cases} \bar{X} = \mu + \frac{\sigma}{\sqrt{n}} \cdot \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \xrightarrow{d} (\mu, \frac{\sigma^2}{n}) \\ S_n = \sum_{i=1}^n X_i = n\bar{X} \xrightarrow{d} (n\mu, n\sigma^2) \end{cases}$$

$$Y = \sum_{i=1}^n X_i \quad \frac{Y - n\mu}{\sigma\sqrt{n}}$$

③ $M_n(t) = M(t) \rightarrow$ 分布收敛定理

$$\text{Proof: } W_n = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}}, \quad W_n \xrightarrow{d} N(0, 1)$$

$$M_n(t) = E[e^{t \cdot \frac{1}{\sqrt{n}} (\sum_{i=1}^n X_i - n\mu)}] = E[\exp((\frac{t}{\sqrt{n}}) (\frac{X_1 - \mu}{\sigma}) \dots \exp((\frac{t}{\sqrt{n}}) \frac{X_n - \mu}{\sigma})]$$

$$= [m(\frac{t}{\sqrt{n}})]^n \quad -h < \frac{t}{\sqrt{n}} < h, \quad m(t) = E[\exp(t \cdot (\frac{X_i - \mu}{\sigma}))] \quad (-h < t < h)$$

$$\text{let } Y = \frac{X_i - \mu}{\sigma} \quad m(0) = 1, \quad m'(0) = 0, \quad m''(0) = 1$$

$$\Rightarrow m(t) = 1 + \frac{m''(0)t^2}{2} \quad t \in [0, t]$$

$$= 1 + \frac{1}{2}t^2 + \frac{[m''(0)-1]t^2}{2}$$

$$n \rightarrow \infty, t_1 \rightarrow 0 \Rightarrow \lim_{n \rightarrow \infty} m_n(t_1) = \lim_{n \rightarrow \infty} (1 + \frac{t_1^2}{2n})^n$$

$$= e^{\frac{1}{2}t_1^2}$$

i.e. the mgf of $N(0, 1)$

8.

Markov's Inequality $P(X \geq c) \leq \frac{E[X]}{c}$ X is a nonnegative random variable and $c > 0$

Chebyshev's Inequality $\forall k \geq 1 \quad P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$ (if $\epsilon = k\sigma \quad P(|X - \mu| \geq \epsilon) \leq \frac{\sigma^2}{\epsilon^2}$)

Proof: let $Y = (X - \mu)^2$, $c = k^2\sigma^2$, $E[Y] = \sigma^2$

$$\Rightarrow P(|X - \mu| \geq k\sigma) = P(Y \geq c) \leq \frac{E[Y]}{c} = \frac{1}{k^2}$$

9. $Z_1, \dots, Z_n$ is said to converge in probability to a random variable Z , often denoted by $Z_n \xrightarrow{P} Z$ [依概率收敛] if for any $\epsilon > 0$, $\lim_{n \rightarrow \infty} P(|Z_n - Z| \geq \epsilon) = 0$

$$\lim_{n \rightarrow \infty} P(|\bar{X} - \mu| \geq \epsilon) = 0$$

Proof: $E[\bar{X}] = \mu$, $\text{Var}[\bar{X}] = \frac{1}{n}\sigma^2$

$$\forall \epsilon > 0 \quad P(|\bar{X} - \mu| \geq \epsilon) \leq \frac{1}{\epsilon^2} \cdot \left(\frac{\sigma}{\sqrt{n}}\right)^2 = \frac{\sigma^2}{n\epsilon^2}$$

$$0 \leq \lim_{n \rightarrow \infty} P(|\bar{X} - \mu| \geq \epsilon) \leq \lim_{n \rightarrow \infty} \frac{\sigma^2}{n\epsilon^2} = 0$$